

## AMS 261: Probability Theory (Fall 2017)

### Homework 4 (due Tuesday 11/28)

1. Consider a sequence  $\{X_n : n = 1, 2, \dots\}$  of  $\overline{\mathbb{R}}$ -valued random variables defined on the same probability space  $(\Omega, \mathcal{F}, P)$ . Assume that the sequence is (pointwise) increasing, that is, for all  $n$  and for each  $\omega \in \Omega$ ,  $X_n(\omega) \leq X_{n+1}(\omega)$ . Moreover, assume that  $E(X_1) > -\infty$ . Denote by  $X$  the pointwise limit of  $\{X_n : n = 1, 2, \dots\}$ , that is, for each  $\omega \in \Omega$ ,  $X(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$ .
  - Prove that  $E(X) = \lim_{n \rightarrow \infty} E(X_n)$ .
2. Let  $\{X_n : n = 1, 2, \dots\}$  be a countable sequence of  $\overline{\mathbb{R}}^+$ -valued random variables defined on a common probability space  $(\Omega, \mathcal{F}, P)$ , and assume that  $E(\sum_{n=1}^{\infty} X_n) < \infty$ .
  - Show that  $E\left(\sum_{n=1}^{\infty} X_n\right) = \sum_{n=1}^{\infty} E(X_n)$ .
3. Let  $\{X_n : n = 1, 2, \dots\}$ ,  $\{Y_n : n = 1, 2, \dots\}$ , and  $\{Z_n : n = 1, 2, \dots\}$  be sequences of  $\mathbb{R}$ -valued random variables (all the random variables are defined on the same probability space). Assume that: (a)  $E(X_n)$  and  $E(Z_n)$  exist for all  $n$  and are finite; (b) each of the three sequences converges almost surely (denote by  $X$ ,  $Y$ , and  $Z$  the respective almost sure limits); (c)  $E(X)$ ,  $E(Y)$ , and  $E(Z)$  exist and are finite; (d)  $X_n \leq Y_n \leq Z_n$  almost surely; (e)  $\lim_{n \rightarrow \infty} E(X_n) = E(X)$ , and  $\lim_{n \rightarrow \infty} E(Z_n) = E(Z)$ .
  - Show that  $\lim_{n \rightarrow \infty} E(Y_n) = E(Y)$ .
4. Let  $\{X_n : n = 1, 2, \dots\}$  be a countable sequence of  $\mathbb{R}$ -valued random variables defined on a common probability space  $(\Omega, \mathcal{F}, P)$ . Assume that there exist finite real constants  $p > 1$  and  $K > 0$  such that  $\sup_n E(|X_n|^p) \leq K$ .
  - Show that  $\{X_n : n = 1, 2, \dots\}$  is uniformly integrable.
5. Let  $X$  be an  $\mathbb{R}$ -valued random variable, defined on probability space  $(\Omega, \mathcal{F}, P)$ , with finite expectation  $\mu = E(X)$  and finite standard deviation  $\sigma = \{\text{Var}(X)\}^{1/2}$ .
  - Prove that for any  $0 \leq z \leq \sigma$ ,
$$P(\{\omega \in \Omega : |X(\omega) - \mu| \geq z\}) \geq \frac{\sigma^4 \{1 - (z/\sigma)^2\}^2}{E(|X - \mu|^4)}.$$
6. Let  $\{X_n : n = 1, 2, \dots\}$  be a sequence of  $\mathbb{R}$ -valued random variables defined on a common probability space  $(\Omega, \mathcal{F}, P)$ . Suppose there exists an  $\mathbb{R}^+$ -valued random variable  $Y$ , defined on  $(\Omega, \mathcal{F}, P)$ , such that  $E(Y) < \infty$  and  $|X_n| \leq Y$ , almost surely, for all  $n$ .
  - Show that  $\{X_n : n = 1, 2, \dots\}$  is uniformly integrable.
7. Consider a countable sequence  $\{X_n : n = 1, 2, \dots\}$  of  $\overline{\mathbb{R}}$ -valued random variables, defined on a common probability space  $(\Omega, \mathcal{F}, P)$ , and an increasing function  $G : [0, \infty) \rightarrow [0, \infty)$ , which satisfies  $\lim_{t \rightarrow \infty} \{t^{-1}G(t)\} = \infty$  and  $0 < \sup_n E\{G(|X_n|)\} < \infty$ .
  - Prove that  $\{X_n : n = 1, 2, \dots\}$  is uniformly integrable.